

NOT ALL HOME DRINKING IS EQUAL

A LATENT CLASS ANALYSIS OF DRINKING PATTERNS AND ALCOHOL CONSUMPTION LEVELS FOLLOWING INITIAL COVID-19 RESTRICTIONS IN AUSTRALIA, NEW ZEALAND, AND THE UNITED KINGDOM



Presented by Dr Cheneal Puljević

ARC DECRA Fellow, NHMRC Centre of Research Excellence on Achieving the Tobacco Endgame, The University of Queensland

Published in *Drug and Alcohol Review* (2023)

Tom R. Strating (1), Cheneal Puljević (1,2), Emma Davies (3), Monica J. Barratt (4,5), Adam Winstock (6,7), Jason Ferris (1)

1. Centre for Health Services Research, The University of Queensland
2. School of Public Health, The University of Queensland
3. The Centre for Psychological Research, Oxford Brookes University
4. Social Equity Research Centre and Digital Ethnography Research Centre, RMIT University
5. National Drug and Alcohol Research Centre, University of New South Wales
6. University College London
7. Global Drug Survey Ltd, London, United Kingdom

BACKGROUND AND AIM

Initial COVID-19 restrictions forced changes in the contexts (e.g., with who and where) within which individuals consumed alcohol.

This study uses data from the Global Drug Survey to explore different profiles of drinking contexts during initial COVID-19 restrictions and their associations with alcohol consumption.




Figure 1: Endorsement profiles for past month alcohol use by latent class for the six-class model




Figure 2: Predictive margins plot showing the number of standard drinks in the last month by sex and age for each class

METHOD

We used latent class analysis (LCA) to explore subgroups of drinking contexts among 4,891 respondents of the Global Drug Survey from the United Kingdom, New Zealand and Australia who reported drinking alcohol in the month prior to data collection (3 May- 21 June 2020). Ten binary LCA indicator variables (shown in Figure 1) were generated from a survey question about last month alcohol settings. Negative binomial regression was used to explore the association between the latent classes and respondents' total number of drinks consumed in the last 30 days (i.e., alcohol consumption).

RESULTS

Figure 1 shows the six distinct classes of individuals who reported drinking in the following contexts: Household (36.0%); Alone (32.3%); Alone & household (17.9%); Gatherings & household (9.5%); Party (3.2%), and Everywhere (1.1%). The Everywhere group was associated with the highest probability of increased alcohol consumption at the time. Figure 2 shows that male respondents and those aged 35 or older were most likely to report increased alcohol consumption.

DISCUSSION

Our findings suggest that drinking contexts, sex and age influenced alcohol consumption during the early stages of the COVID-19 pandemic. These findings highlight a need for improved policy targeting risky drinking in home settings. Further research should explore whether COVID-19-induced shifts in alcohol use persist as restrictions are lifted.

TAKE HOME MESSAGE

Drinking contexts, sex, and age influenced alcohol consumption during the early stages of the COVID-19 pandemic

