Anabolic Steroid Use: A Looming Public Health Problem?

Harrison G. Pope, Jr., M.D. November 14, 2017

McLean Hospital / Harvard Medical School Belmont, MA, USA



Anabolic-Androgenic Steroids (AAS):

Testosterone: The natural male hormone

Synthetic hormones similar to testosterone:

- Nandrolone (Deca-Durabolin)
- Methenolone (Anadrol)
- Methandienone (Dianabol)
- Stanozolol (Winstrol)

History of AAS:

- 1935: Testosterone first isolated by David et al.
- 1940's: Many synthetic AAS developed
- 1950-1980: AAS spreads through elite athletics
- 1980's: AAS use spreads to non-athletes
- 1990-2010: AAS use becomes widespread
- among young men in Western countries

Thus, most illicit AAS users are young, rankand-file gym clients who are NOT competitive athletes.



AAS Use Among American Men

Myths About AAS Use

Most AAS users are competitive athletes.

AAS use is common in adolescents.

Girls and women often use AAS.

Why do People Use AAS?:

I. Highly Effective Drugs

II. Supportive Social Climate

III. Individuals with Body-Image Concerns



































Western Mythological Tradition:

Bernadino Genga's drawing of the Farnese statue of Herakles (1691)



Scandinavian Mythological Tradition:

Johann Heinrich Füssli's Der Kampf des Thor mit der Schlange des Midgard (1788)



Japanese Mythological Tradition:

Eitaku Kobayashi's *Izanagi and Izanami* (1885)



Japanese Mythological Tradition:

Unknown sculptor: Seated Male Deity (c. 800)



III. Individuals With Body-Image Concerns



Time to onset of AAS use for men with high vs. low adolescent body-image concerns









What Should We Be Worrying About?

Cardiovascular effects
Persistent hypogonadism
Psychiatric disorders
Neurotoxicity

Dilated Cardiomyopathy









11/15/2017





Coronary CT Angiography





CT Angiography: Noncalcified plaque in right



Persistent Hypogonadism

- In males, exogenous AAS suppress the hypothalamicpituitary-testicular axis.
- If a man stops AAS after a prolonged "cycle," he will likely be hypogonadal.
- Hypogonadism may resolve within weeks, but can persist more than a year.



Psychiatric Disorders

 Hypomanic or manic episodes during AAS exposure

 Major depression during AAS withdrawal







NORWAY, 2011

Major Depressive Episodes

- Usually associated with hypogonadism during AAS withdrawal
- Affects only a minority of users
- In rare cases may lead to suicide



Continued Blebbing

Study Participants

- Long-term AAS users (at least two years of total lifetime AAS use), N = 22
- Short-term AAS users (less than two years of total lifetime AAS use), N = 9
- Experienced weightlifters reporting no AAS use, N = 13





Pattern Recognition Memory, Errors

Paired Associates Learning, Errors





Paired Associates Learning, Errors





Magnetic Resonance Spectroscopy



A Looming Public Health Problem?

 Widespread illicit AAS use did not begin until about 1980.

Therefore the leading edge of the population of illicit AAS users – those who first tried steroids as youths in the 1980's – is only now reaching middle age.

A Looming Public Health Problem?

Imagine that widespread cigarette smoking did not arise until 1980, and that the great majority of cigarette smokers were still under age 50 today...

 An analogous situation might presently exist for AAS.