



Reinfection following successful HCV DAA therapy among people with recent injecting drug use: the SIMPLIFY and D3FEAT studies

Evan Cunningham, Grebely J, Dalgard O, Hajarizadeh B, Conway B, Powis J, Bruneau J, Feld JJ, Read P, Cooper C, Amin J, Bruggmann P, Lacombe K, Stedman C, Hellard ME, Marks P, Dunlop A, Quiene S, Moriggia A, Applegate TL, Litwin AH, Matthews GV, and Dore GJ on behalf of the SIMPLIFY and D3FEAT Study Groups

21 September, 2018



Disclosures

Nothing to disclose

Background/rationale

- There is a significant burden of hepatitis C virus infection among people who inject drugs globally¹
- Treatment has been shown to be safe and effective in people who inject drugs
 - 94% SVR in SIMPLIFY and 91% in D3FEAT
- Reinfection following therapy has been one of the major concerns around scale up of HCV DAA treatment among people who inject drugs
- There is limited data on reinfection following HCV DAA therapy among people with ongoing injecting risk behaviours

¹Grebely et al, 2018, Addiction

Aims

1. Assess the incidence of HCV reinfection, including stratification by key risk behaviours.
2. Investigate predictors of time to HCV reinfection

SIMPLIFY and D3FEAT study Design

- Investigator-initiated, Kirby/UNSW sponsored, international open-label trials
- 25 sites, 8 countries
- Study recruitment conducted through a network of drug and alcohol clinics, hospital clinics, and community clinics
- Participants enrolled between April 2016 and February 2017



Study design and participant eligibility

- DAA treatment-naïve patients with GT1-6 chronic HCV infection (F0-4)
- Treated with sofosbuvir and velpatasvir (SIMPLIFY; n=97) or PrOD±RBV (D3FEAT; n=82)
- People with recent injecting drug use (past six months; SIMPLIFY) or people with either recent injecting drug use or currently on OST (D3FEAT)
- Participants with HIV and decompensated liver disease excluded



Reinfection

- Measured every 6 months following SVR24
 - Tested for HCV RNA
 - Complete follow-up questionnaire
- Reinfection assessed from end of treatment
 - Distinguished from relapse using viral sequencing



Study outcome and statistical analysis

Reinfection

- Quantifiable HCV RNA following HCV DAA therapy
 - Distinguished from HCV relapse using viral sequencing
- Rate calculated using person-time (cases per 100 person-years)

Participant characteristics

Characteristic	DAA treatment (12 weeks) N = 179
Female, n (%)	48 (27%)
Age, median years (25%, 75%)	48 (42, 54)
Current opioid substitution therapy, n (%)	108 (60%)
Injecting at EOT	
Any injecting drug use	97 (54%)
Daily or greater injecting	34 (19%)
Injecting following EOT	
Any injecting drug use	124 (69%)
Daily or greater injecting	52 (29%)
Heroin injecting	82 (46%)
Methamphetamine injecting	52 (29%)
Other opioid injecting	43 (24%)
Cocaine injecting	34 (19%)


Reinfections

- Overall there were 9 cases of viral recurrence including 6 reinfections and 3 relapses


Risk behaviours post EOT

ID	Sex	Age	Country	Injecting drug use	Highest frequency	Main drug	Sharing needles
16	Male	36	Canada	Yes	≥daily	Morphine	No
27	Male	55	Canada	Yes	≥daily	Morphine	No
93	Male	41	Australia	Yes	≥daily	Heroin	No
98	Male	24	New Zealand	Yes	≥daily	Methamphetamine	No
59	Male	32	Switzerland	Yes	≥daily	Cocaine	Yes
87	Female	28	New Zealand	Yes	<daily >monthly	Morphine	Yes


Incidence of reinfection



Incidence of reinfection



Incidence of reinfection



Discussion

- Reinfection following successful HCV DAA therapy does occur
- All observed reinfections occurred among people with ongoing injecting after ETR
- Higher incidence of reinfection in those with more frequent injecting
- The incidence of reinfection is consistent with previously reported rates of reinfection in the interferon era
- DAA treatment has the potential to be used as an opportunity to encourage safe injection practices and uptake of harm reduction

Acknowledgements



SIMPLIFY and D3FEAT study participants

Study coordination staff: Sophie, Amanda, Pip, Ecaterina, Mahshid



SIMPLIFY study group

D3FEAT study group

Protocol Steering Committee – Gregory Dore (Chair, UNSW Sydney, Sydney, Australia), Philip Bruggmann (Arud Centres for Addiction Medicine, Zurich, Switzerland), Jason Grebely (UNSW Sydney, Sydney, Australia), Philipp Marks (UNSW Sydney, Sydney, Australia), Julie Bruneau (Centre Hospitalier de l'Université de Montréal, Canada), Tracy Swan (Médecins Sans Frontières, New York, United States), Olav Dalgard (Akershus University Hospital, Oslo, Norway), Jude Byrne (Australian Injecting & Illicit Drug Users League), Melanie Lacalamita (Poliklinik für Infektiologie, Inselspital, Bern, Switzerland) and Adrian Dunlop (Newcastle Pharmacotherapy Service, Newcastle, Australia).

Coordinating Centre – Sophie Quiene (Study Co-ordinator), Evan Cunningham (PhD Student), Behzad Hajarizadeh (Associate Lecturer), Gregory Dore (co-Principal Investigator), Jason Grebely (co-Principal Investigator), Pip Marks (Clinical Trials Manager), Ineke Shaw (Systems Manager), Sharmila Siriragavan (Data Manager) and Janaki Amin (Statistician).

Site Principal Investigators – Philip Bruggmann (Arud Centres for Addiction Medicine, Zurich, Switzerland), Julie Bruneau (Centre Hospitalier de l'Université de Montréal, Montréal, Canada), Brian Conway (Vancouver Infectious Diseases Center, Vancouver, Canada), Olav Dalgard (Akershus University Hospital, Oslo, Norway), Gail Matthews (St Vincent's Hospital, Sydney Australia), Adrian Dunlop (Newcastle Pharmacotherapy Service, Newcastle, Australia), Margaret Hellard (The Alfred Hospital, Melbourne, Australia), Jeff Powis (South Riverdale Community Health Centre, Toronto, Canada), David Shaw (Royal Adelaide Hospital, Adelaide, Australia), Maria Christine Thumheer (Poliklinik für Infektiologie, Inselspital, Bern, Switzerland), Martin Weitman (Nepean Hospital, Penrith, Australia), Ian Kronborg (Footscray Hospital, Footscray, Australia), Curtis Cooper (The Ottawa Hospital, Ottawa, Canada), Jordan Feld (Toronto General Hospital, Toronto, Canada), Christopher Fraser (Coolaid Community Health Centre, Victoria, Canada), Alain Litwin (Montefiore Medical Centre, New York, United States), John Dillon (Ninewells Hospital, Dundee, United Kingdom), Ed Gane (Auckland Hospital, Auckland, New Zealand), Phillip Read (Kirketon Road Centre, Sydney, Australia).

Site Co-ordinators – Jessica Andreassen, Ingunn Melkeraaen and Merete Moen Tollefsen (Akershus University Hospital, Oslo, Norway), Catherine Ferguson (Royal Adelaide Hospital, Adelaide, Australia), Nargis Abram and Vincenzo Fragomeli (Nepean Hospital, Penrith, Australia), Susan Hazelwood and Michelle Hall (Newcastle Pharmacotherapy Service, Newcastle, Australia), Tina Horschik (Arud Centres for Addiction Medicine, Zurich, Switzerland), Marie-Claire Chayer and Barbara Kotsoros (Centre Hospitalier de l'Université de Montréal, Montréal, Canada), Melanie Lacalamita (Poliklinik für Infektiologie, Inselspital, Bern, Switzerland), Kate Mason (South Riverdale Community Health Centre, Toronto, Canada), Alison Sevehon (St Vincent's Hospital, Sydney, Australia), Hannah Pagarigan (Vancouver Infectious Diseases Center, Vancouver, Canada), Michelle Hagenauer (The Alfred Hospital, Melbourne, Australia), Rachel Liddle (Footscray Hospital, Footscray, Australia), Miriam Muir and Jessica Millroy (The Ottawa Hospital, Ottawa, Canada), Diana Kaznowski and Lily Zou (Toronto General Hospital, Toronto, Canada), Rozalyn Milne (Coolaid Community Health Centre, Victoria, Canada), Linda Agyemang and Hiral Patel (Montefiore Medical Centre, New York, United States), Shirley Clearly and Linda Johnston (Ninewells Hospital, Dundee, United Kingdom), Victoria Oliver (Auckland Hospital, Auckland, New Zealand), Rebecca Lothian and Rosemary Gilliver (Kirketon Road Centre, Sydney, Australia).