



Up to
12
HOURS
Of Clinically Proven
Pain Relief^{1,2}



**EXPANDED
LABEL INDICATION**

Can be used for targeted relief of localised muscle pain or mild OA of the knees and fingers³

The POWER OF DICLOFENAC for up to 12 hours relief of musculoskeletal pain and inflammation¹⁻⁶

Voltaren Pain Relief Gel 12 Hourly can be used for up to:³

- 2 weeks to relieve pain in acute soft tissue injury including sprains, strains, sporting injuries and tendonitis.
- 3 weeks to relieve pain from mild and localised forms of osteoarthritis of the knees and fingers.



SPORT & OVERUSE INJURIES

Generally caused by repetitive movements or moving incorrectly and overloading a tendon, e.g., elbow tendonitis⁷



OSTEOARTHRITIS PAIN

Occurs when cartilage becomes damaged, with hips, knees and hands commonly affected^{8,9}



SPRAINS

Overstretching of ligaments, e.g., ankle sprain^{10,11}




STRAINS

Damage to muscles or tendons by oversteering, e.g., hamstring strain^{10,11}


A proven anti-inflammatory effect with pain relief targeted to the source^{4-6,12}

Diclofenac, like most NSAIDs, inhibits the activity of the enzyme cyclooxygenase, reducing the production of prostaglandins and therefore inflammation.^{12,13}



Voltaren Pain Relief Gel 12-Hourly has a **hydrogel phase** and a **lipid phase** containing diclofenac^{14,15}

••• = diclofenac • = lipid



The epidermis acts as a **reservoir for sustained release of diclofenac** into underlying tissues^{14,15}



Hydrogels in the gel base allow water evaporation, providing a **cooling and soothing effect on the skin**^{14,15}



A **permeation enhancer** allows up to **3X greater skin penetration** to act at the source of pain^{16*}



On skin contact, **lipids** release diclofenac enabling **penetration through the layers**^{14,15}




Up to **12 hours pain relief** per application³

*Voltaren Pain Relief Gel 12 Hourly vs Voltaren Pain Relief Gel, *in vitro* study.¹⁶

Expert consensus guidelines support topical NSAIDs for musculoskeletal pain¹⁷⁻¹⁹

ACUTE MUSCULOSKELETAL PAIN



First-line therapy:

Topical NSAIDs, with or without menthol gel, to reduce or relieve symptoms, including pain; improve physical function; and improve treatment satisfaction

Strong recommendation with moderate-certainty evidence.

*American College of Physicians; American Association of Family Medicine*¹⁷

OSTEOARTHRITIS



Strong recommendation:

Topical NSAIDs for knee osteoarthritis in patients with no comorbidities, GI or CV comorbidities or frailty. *Osteoarthritis Research Society International*¹⁸

Primary option:

Topical NSAIDs for knee osteoarthritis in patients over 16 years, to reduce symptoms and support non-pharmacological management. Benefit may accrue for other osteoarthritis-affected joints. *NICE Guidelines*¹⁹

Recommend Voltaren Pain Relief Gel 12 hourly for your patients



Up to 12 hours of clinically proven pain relief^{1,2}



Suitable for acute soft tissue injuries such as sports injuries³



Suitable for mild osteoarthritis of knees and fingers³



Targets the site of pain and inflammation^{14,15,20}



3X greater skin penetration^{16*}



Low systemic absorption⁶



Up to 3 weeks consecutive use^{3†}



Easy to open cap

*Voltaren Pain Relief Gel 12 Hourly vs Voltaren Pain Relief Gel, *in vitro* study.¹⁶ †For mild and localised forms of OA of the knees and fingers.³

Voltaren Pain Relief Gel 12 Hourly should not be used by:³



Children under 12 years



Women in the last 3 months of pregnancy



In people with an allergy to NSAIDs

Voltaren Pain Relief Gel 12 Hourly should be used with caution by:³



Women planning to become pregnant

Do not use:³



- For more than 2 weeks for acute soft tissue injury
- For more than 3 weeks for mild OA of the knees and fingers

NEW LOOK VOLTAREN RANGE

With options to suit your patient's needs

TOPICAL PAIN RELIEF

ORAL PAIN RELIEF



VOLTAREN PAIN RELIEF GEL 12 HOURLY

For long-lasting and targeted relief of localised muscle pain or mild OA of the knees and fingers³



VOLTAREN PAIN RELIEF GEL

For targeted relief of localised muscle pain or mild OA of the knees and fingers²¹



VOLTAREN RAPID EXTRA STRENGTH

Fast relief of strong pain that lasts up to 8 hours²²⁻²⁵



VOLTAREN RAPID

Relief of acute pain including headache and cold/flu symptoms²⁶

Always read the label and follow the directions for use. Incorrect use could be harmful. Voltaren Rapid Extra Strength: Ask your pharmacist about this product. Incorrect use could be harmful.

Voltaren Pain Relief Gel 12 Hourly contains diclofenac diethylamine 23.2 mg/g. Indications and dosage: For (up to 2 weeks) relief of pain from acute soft tissue injury including sprains, strains, sporting injuries and tendonitis; for (up to 3 weeks) relief of pain from mild and localised forms of osteoarthritis of the knees and fingers. Adults and children ≥ 12 years: Apply to affected area up to 2 times a day. Contraindications: Hypersensitivity to any ingredient; asthma attacks, urticaria or rhinitis precipitated by aspirin or other NSAIDs; in the last 3 months of pregnancy; in children under 12 years. Adverse reactions (common): Rash, eczema, dermatitis, pruritus. For less common adverse reactions see Pack insert. **Voltaren Pain Relief Gel** contains diclofenac diethylamine 11.6 mg/g. Indications and dosage: Adults and children ≥ 12 years: For (up to 2 weeks) relief of pain from acute soft tissue injury including sprains, strains, sporting injuries and tendonitis; for (up to 3 weeks) relief of pain from mild and localised forms of osteoarthritis of the knees and fingers. Apply to affected area up to 4 times daily. Contraindications: Hypersensitivity to any ingredient; in asthma attacks, urticaria or rhinitis precipitated by aspirin or other NSAIDs; in the last 3 months of pregnancy; in children under 12 years. Adverse reactions (common): Rash, eczema, dermatitis, pruritus. For less common adverse reactions see Pack insert. **Voltaren Rapid Extra Strength: Australia: Please review Product Information (PI) before recommending Voltaren Rapid Extra Strength The full PI is available from Haleon Australia on request (FREECALL 1800 028 533).** Voltaren Rapid Extra Strength contains diclofenac potassium 25 mg. Use: For the temporary relief of acute pain where inflammation is present, such as sprains and strains, back and joint pains, migraine headaches and period pain. Adults: 2 tablets initially with liquid, then 1–2 tablets every 8 hours as necessary. Children 14–17 years: Take 1–2 tablets initially with liquid, then 1 tablet every 8 hours as necessary. (Maximum 8 tablets in 24 hours). Contraindications: Hypersensitivity to any ingredient; in asthma attacks, urticaria or rhinitis precipitated by aspirin or other NSAIDs; patients with previous myocardial infarction within the last 6–12 months; severe cardiac failure; hepatic failure; renal failure; active gastric or intestinal ulcer, bleeding or perforation; last trimester of pregnancy; children under 14 years. **Voltaren Rapid** contains diclofenac potassium 12.5 mg. Use: For the temporary relief of painful conditions such as headache, dental pain, period pain, rheumatic and muscular pain, backache. Temporary relief of symptoms of colds and flu (including aches and pains, sore throat pain). Reduces fever. Adults and children ≥ 14 years: Take 2 tablets initially with liquid, then 1–2 tablets every 4–6 hours if necessary (max 6 tablets per day). For further information, please contact 1800 028 533.

References: 1. Derry S, et al. Cochrane Database Syst Rev 2015;6:CD007042. 2. Predel HG, et al. Med Sci Sports Exer 2012;44(9):1629–36. 3. Voltaren Pain Relief Gel 12 Hourly package insert. 4. Derry S, et al. Cochrane Database Syst Rev 2016;4:CD007400. 5. Predel HG, et al. BMC Musculoskeletal Disord 2013;14:250. 6. Derry S et al. Cochrane Database Syst Rev 2017;5:CD008609. 7. Mayo Clinic. Tendinitis. Available at: www.mayoclinic.org/diseases-conditions/tendinitis/symptoms-causes/syc-20378243. Accessed December 2024. 8. Arthritis Foundation. What is Osteoarthritis? Available at: www.arthritis.org/diseases/osteoarthritis. Accessed December 2024. 9. Cushnaghan J, Dieppe P. Ann Rheum Dis 1991;50:8–13. 10. Safer Care Victoria. Soft tissue injuries (Sprains and strains). Available at: www.safercare.vic.gov.au/sites/default/files/2019-07/Soft%20tissue%20injuries.pdf. Accessed December 2024. 11. Sports Medicine Australia. Soft Tissue Injuries. Available at: sma.org.au/resources-advice/injury-fact-sheets/soft-tissue-injuries/. Accessed December 2024. 12. Zacher J, et al. Curr Med Res Opin 2008;24:925–50. 13. Ricciotti E, FitzGerald G. Arterioscler Thromb Vasc Biol 2011;31(5):986–1000. 14. Haleon Data on File. Diclofenac skin penetration by vibrational spectroscopy and imaging. 15. Gondolph-Zink B et al. Akt Rheumatol 1996;21:298–304. 16. Quartarone G, Hasler-Nguyen N. Arthritis Rheumatol 2014;66:S976. 17. Qaseem A, et al. Ann Int Med 2020;173:739–48. 18. Bannuru RR, et al. Osteoarthritis Cartilage 2019;27(11):1578–89. 19. NICE. Osteoarthritis in over 16s: diagnosis and management. Published 19 October 2022. Available at: www.nice.org.uk/guidance/ng226. Accessed December 2024. 20. Brune K. Curr Med Res Opin 2007;23(12):2985–95. 21. Voltaren Pain Relief Gel package insert. 22. Bakshi R et al. Curr Med Res 1992;52(3):435–42. 23. Kintigh JW. Today Ther Trends 1995;Suppl 1:47–61. 24. Sumatriptan Migraine Study Group. Cephalgia 1999;19:232–40. 25. Voltaren Rapid Extra Strength Australian approved Product Information. 26. Voltaren Rapid package insert.

Trademarks are owned by or licensed to the Haleon group. ©2025 Haleon or its licensor. Sydney, NSW Australia & Auckland, New Zealand. April 2025. PM-AU-VOLT-25-00089. HALEOY0165.

HALEON