



Sanofi Vaccines for Travellers



DISEASE	VACCINE	AGE LIMIT	DOSE	INITIAL SCHEDULE	NOTES
Hepatitis A	AVAXIM® Hepatitis A vaccine (inactivated, adsorbed)	≥2 years	0.5 mL	Single dose ¹	Booster at 6–36 months ¹
Typhoid fever	TYPHIM Vi® <i>Salmonella typhi</i> Vi polysaccharide vaccine	≥2 years	0.5 mL	Single dose ²	Revaccination recommended every 3 years in patients who remain at risk. ²
Rabies	VERORAB® Inactivated Rabies Virus	Any age	0.5 mL (IM) or 0.1 mL (ID)	Pre-exposure: ³ - Conventional regimen Day 0, 7 and 21 or 28 - One week regimen ^{4,5} Day 0, 7	*This regimen should not be used for immunocompromised individuals. ³ For ID administration, give two doses at both day 0 and day 7. ³ Refer to Verorab Australian Product Information for information regarding one-week regimen and post-exposure dosing. ³ For information regarding boosters refer to the Australian Immunisation Handbook.
Yellow fever	STAMARIL® Yellow Fever Vaccine (live)	≥9 months	0.5 mL	Single dose ⁴	Refer to Australian Immunisation Handbook for booster recommendations; refer to PI for patients older than 60 years. ^{4,10}
Meningococcal meningitis	MENQUADFI® Meningococcal (Groups A, C, Y & W-135) Polysaccharide Tetanus Toxoid Conjugate Vaccine	≥12 months	0.5 mL	Single dose ⁵	Refer to MenQuadfi Australian Product Information for information regarding booster vaccination in adolescents & adults who have previously been primed with meningococcal vaccine at least 4 years prior. ⁵
Diphtheria, Tetanus, Pertussis	ADACEL® Pertussis Vaccine-Acellular Combined with Diphtheria and Tetanus Toxoids (Adsorbed) ¹	≥10 years	0.5 mL	Single dose ⁶	Use to revaccinate following primary vaccination. Can be used for repeat booster vaccination at 5–10 year intervals. ⁶
Diphtheria, Tetanus, Pertussis, Poliomyelitis	ADACEL® POLIO Pertussis Vaccine-Acellular Diphtheria and Tetanus Toxoids (Adsorbed) Combined with inactivated Poliovirus Type 1, 2 and 3 (Vero cell)	≥4 years	0.5 mL	Single dose ⁷	Use to revaccinate following primary vaccination. Can be used for repeat booster vaccination at 5–10 year intervals. ⁷
Poliomyelitis	IPOL® Inactivated Poliomyelitis Vaccine (Vero)	≥2 months	0.5 mL	3 doses 8 weeks apart ⁸	Children: booster dose recommended at 4 years of age for patients who have already received 3-dose primary series. Adults: booster dose 10-yearly for travellers and those at continued risk. ⁸

Influenza Vaccines

Influenza Types A & B (Quadrivalent)	FLUQUADRI™ Inactivated Quadrivalent Influenza Vaccine, Split Virion (Influenza virus haemagglutinin)	≥6 months	0.5 mL	Single dose ⁹	As different strains circulate from year to year, annual vaccination with current formulation is necessary. ¹⁰ ⁹ Children aged 6 months to <9 years receiving influenza vaccine for the first time, should receive two doses 4 weeks apart. ^{9,10} Refer to the Fluquadri Australian Product Information and the Australian Immunisation Handbook for further information. ^{9,10}
	FLUZONE® HIGH-DOSE QUADRIVALENT Inactivated quadrivalent influenza vaccine, split virion (Influenza virus haemagglutinin)	≥60 years	0.7 mL	Single dose ¹¹	As different strains circulate from year to year, annual vaccination with the current formulation is necessary. ¹⁰ Contains 60µg HA of each of the four strains recommended for the upcoming influenza season. ¹¹ Fluzone High-Dose Quadrivalent should be given in accordance with the national recommendation as per the current Immunisation Handbook. Not available on the NIP. Refer to the Fluzone High-Dose Quadrivalent Australian Product Information and the Australian Immunisation Handbook for further information. ^{10,11}

PBS Information: These products are not listed on the PBS. FluQuadri, Adacel & IPOL are listed on the National Immunisation Program (NIP). Refer to NIP schedule and State or Territory Health Departments for details. All other vaccines are not listed on the NIP.

Before prescribing please review Product Information available on request from Sanofi or online <https://medsinfo.com.au/>

Abbreviations: HA: haemagglutinin; ID: intradermal; IM: intramuscular; NIP: National Immunisation Program; PI: product information.

References: 1. AVAXIM® Australian Product Information. 2. TYPHIM Vi® Australian Product Information. 3. VERORAB® Australian Product Information. 4. STAMARIL® Australian Product Information. 5. MENQUADFI® Australian Product Information. 6. ADACEL® Australian Product Information. 7. ADACEL® POLIO Australian Product Information. 8. IPOL® Australian Product Information. 9. FLUQUADRI™ Australian Product Information. 10. Australian Immunisation Handbook. <https://immunisationhandbook.health.gov.au/vaccine-preventable-diseases> (Accessed November 2021). 11. Fluzone® High-Dose Quadrivalent Australian Product Information.

sanofi-aventis australia Pty Ltd. ABN 31 008 558 807. Trading as SANOFI. 12–24 Talavera Rd, Macquarie Park NSW 2113. Customer Service 1800 829 468. MAT-AU-2302796-V1.0. February 2024.

sanofi