

Introduction

Several nations have committed to reaching net-zero emissions to mitigate the disastrous effects of climate change. India, too, has set up a timeline to reach net zero by 2070, announced at COP26. As a developing nation, India will need to deploy CCS to meet its goals while taking care of energy security for the growing population sustainably. Moreover, geologic formations in India offer an enormous potential for carbon sequestration [Vishal, et al., 2021]. Storage of CO₂ in depleted oil and gas fields has been identified as one of the primary pathways to advance CCS at a large scale in India [NITI Aayog, 2022]. It offers dual benefits of abatement CO₂ abatement and extraction of unrecoverable hydrocarbons simultaneously through CO₂-enhanced oil recovery (EOR). This technology can help India keep its net-zero commitment while ensuring sustainable economic progress.

However, no projects are currently running for CO₂-EOR in India. EOR is primarily carried out through water flooding in most fields primarily to maintain reservoir pressure. Nonetheless, there are a few pilot projects planned. Gandhar field in the Cambay basin in western India has been selected by the Government of India as one of the primary prospects for implementing CO₂ EOR in the country [Mishra et al., 2023]. Feasibility studies are also being carried out to screen potential sites in other basins [MoPNG, 2022].

In the Upper Assam Basin, a mature hydrocarbon province, CO₂-EOR demonstrates a 10% incremental recovery advantage over water flooding while enabling subsurface CO₂ storage through residual trapping and solubility mechanisms [Dutta et al., 2024]. Previous studies show that systems utilizing anthropogenic CO₂ from natural gas processing show 15-30% lower emissions per barrel of produced crude compared to conventional water flooding, primarily through avoided venting and displacement of higher-emission power generation [Jensen et al., 2018].

Before implementing a CCS project, it is vital to assess the net environmental benefit of CO₂ capture and its utilization for CO₂-EOR. Life Cycle Assessment (LCA) is a widely used methodology for analyzing these technologies for their net emissions and identifying appropriate improvements to make the process more effective. Due to the relatively low TRL of CO₂ EOR in India [Vishal, et al., 2021], no such study has been conducted incorporating the storage aspects.

Most of the current LCA models developed for CCUS are based on studies carried out in the United states and Europe. However, given the complexities involved in the Indian scenario, they cannot suitably provide accurate results when implemented for potential CCS projects in the country. There is a need for assessment of life cycle emissions based on Indian specific conditions and data. Furthermore, the feasibility and potential benefit of CO₂ EOR over water flooding in Indian basins as a primary oil recovery process must be analyzed. In this study, we aim to assess the avoided CO₂ in the process and its benefit in reducing greenhouse gas (GHG) emissions.

Method

We develop an LCA model for CO₂ and waterflood EOR based on India-specific technologies and parameters. Our study aims to understand and compare the system boundary for CO₂- with water flood EOR using lifecycle assessment process. We carry out detailed gate-to-gate LCA for both processes and their various sub-units to derive the potential benefit in terms of abatement of GHG emissions. We have selected an oil field in western India to assess the environmental impact of CO₂- as well as waterflood EOR. This provides a much-needed comparison of the various options available to make informed decisions.

Results

The key LCA parameters influencing the total emissions include the crude recovery ratio and CO₂ utilization efficiency. Higher CO₂-oil recovery ratios reduce emissions intensity when using captured industrial CO₂, but increase it when employing geologically sourced CO₂ due to lost displacement credits for clean electricity. For water flooding, environmental impacts shift to water resource depletion risks and produced water management challenges, particularly in flood-prone regions.

EOR potential in Indian basins is significant. However, full lifecycle accounting must consider methane leakage risks during EOR operations and long-term monitoring costs. The findings underscore CO₂-EOR's dual role in India's energy transition – extending productive lifespans of mature fields while providing pilot project sites for CCS technologies critical for decarbonizing heavy industries.

Conclusions

India faces some unique challenges in CCS compared to other developed countries. To move from waterflood to CO₂ EOR as the primary recovery technology for sustainable development, large-scale projects need to be identified. Our study highlights the role of CO₂ EOR in decarbonization of hard-to-abate sectors, especially the oil and gas industry. Our gate-to-gate lifecycle assessment shows the environmental benefits of CO₂ EOR.

However, the oil-producing basins are distant from India's coal-producing and utilization regions. The source of CO₂ in western India will most likely be either industrial hubs or oil refineries. In contrast, most of the CO₂ will be sourced from the cement industry or thermal power plants in eastern and central India. Currently, networks of pipelines across the vast Indian subcontinent for CO₂ transportation are almost non-existent. The transportation of CO₂ and refined oil products is unique to India and has to be analyzed accordingly for its environmental impact. Moreover, it will also have to be analyzed in conjunction with the sink availability as most of the CO₂-EOR will be carried out in the western and north-eastern parts of India.

References

- Dutta, R., Kundu, G., Mousa, S., Mirkalaei, M., Chakraborty, R., Yomdo, S., and Mandal, A. [2024]. Evaluation of Potential of CO₂-Enhanced Oil Recovery (EOR) and Assessment of Capacity for Geological Storage in a Mature Oil Reservoir within Upper Assam Basin, India. *Energy & Fuels*. <https://doi.org/10.1021/ACS.ENERGYFUELS.4C02384>
- Jensen, M. D., Azzolina, N. D., Schlasner, S. M., Hamling, J. A., Ayash, S. C., and Gorecki, C. D. [2018]. A screening-level life cycle greenhouse gas analysis of CO₂ enhanced oil recovery with CO₂ sourced from the Shute Creek natural gas-processing facility. *International Journal of Greenhouse Gas Control*, 78, 236–243. <https://doi.org/10.1016/j.ijggc.2018.08.010>
- Mishra, G. K., Meena, R. K., Mitra, S., Sahu, S. K., and Aman, B. S. [2023]. Designing an Integrated CO₂-EOR as CCUS cum Blue Hydrogen Demonstration Project: A step towards Net-Zero ONGC-2038. *SPG 14th Biennial International Conference & Exposition*. Kochi: SPG India.
- MoPNG. [2022]. *Draft 2030 Roadmap for Carbon Capture Utilization and Storage for Upstream E&P companies*. Retrieved from <https://mopng.gov.in/en/page/33>
- NITI Aayog. [2022]. Carbon Capture, Utilization and Storage (CCUS) - Policy Framework and its Deployment Mechanism in India. In *NITI Aayog* (Vol. 235). Retrieved from <https://www.niti.gov.in/sites/default/files/2022-12/CCUS-Report.pdf>
- Vishal, V., Chandra, D., Singh, U., and Verma, Y. [2021]. Understanding initial opportunities and key challenges for CCUS deployment in India at scale. *Resources, Conservation and Recycling*, 175(July), 105829. <https://doi.org/10.1016/j.resconrec.2021.105829>
- Vishal, V., Verma, Y., Chandra, D., and Ashok, D. [2021]. A systematic capacity assessment and classification of geologic CO₂ storage systems in India. *International Journal of Greenhouse Gas Control*, 111(February), 103458. <https://doi.org/10.1016/j.ijggc.2021.103458>